


Questions to Ask Your Healthcare Provider

 **Prepopik**[®]
(sodium picosulfate, magnesium oxide, and anhydrous citric acid) for oral solution
10 mg/3.5 g/12 g per packet

- What are the risk factors for colorectal cancer? _____

- Is it time for me to get a colonoscopy? _____

- How do I have to prepare for my colonoscopy? _____

- Is PREPOPIK[®] right for me? _____

- Can I continue to take my medications during the prep? _____

- Who will actually perform the colonoscopy? _____

- How far in advance do I need to arrive at the facility? _____

- How long will the colonoscopy take? _____

- How will I be sedated? What sedative options are available? _____

- Additional Questions/Notes: _____

Indication and Important Safety Information

PREPOPIK[®] is a prescription medicine used by adults to clean the colon before a colonoscopy. PREPOPIK[®] cleans your colon by causing you to have diarrhea. Cleaning your colon helps your healthcare provider see the inside of your colon more clearly during your colonoscopy.

- Do not take PREPOPIK[®] if your healthcare provider has told you that you have serious kidney problems, a blockage in your intestine (bowel obstruction), an opening in the wall of your stomach or intestines (bowel perforation), a very dilated intestine (toxic megacolon), problems with the emptying of food and fluid from your stomach (gastric retention), or an allergy to any of the ingredients in PREPOPIK[®]
 - Tell your healthcare provider right away if you have any of these symptoms resulting from a loss of too much body fluid (dehydration) while taking PREPOPIK[®]: vomiting that prevents you from keeping down the additional prescribed amounts of clear liquids that you must drink after taking PREPOPIK[®], dizziness, urinating less often than normal, or headache
 - PREPOPIK[®] and other bowel preparations can cause serious side effects, including serious loss of body fluid (dehydration) and changes in blood salts (electrolytes) in your blood. These changes can cause abnormal heartbeats that may result in death, seizures (this can happen even if you have never had a seizure), or kidney problems. Your chance of having fluid loss and changes in blood salts with PREPOPIK[®] is higher if you have heart problems, have kidney problems, or take water pills or non-steroidal anti-inflammatory drugs (NSAIDs)
 - The most common side effects of PREPOPIK[®] include nausea, headache, and vomiting. These are not all the possible side effects of PREPOPIK[®]
- You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please see accompanying full Product Information including Medication Guide.

QUESTIONS?

For additional resources, visit PREPOPIK.com. For customer service inquiries, call **1-888-FERRING** (1-888-337-7464).



PREPOPIK[®] is a registered trademark of Ferring B.V.
© 2017 Ferring B.V. All rights reserved.
PK/860/2015/US(1)